



Slovak Society of Chemical Engineering  
Institute of Chemical and Environmental Engineering  
Slovak University of Technology in Bratislava

## PROCEEDINGS

51<sup>st</sup> International Conference of the Slovak Society of Chemical Engineering SSCHE 2025

Hotel DRUŽBA  
Jasná, Demänovská Dolina, Slovakia  
May 27 - 30, 2025

Editors: Assoc. Prof. Mário Mihaľ

ISBN: 978-80-8208-158-2, EAN: 9788082081582

Published by the Faculty of Chemical and Food Technology Slovak Technical University in Bratislava in Slovak Chemistry Library for the Institute of Chemical and Environmental Engineering; Radlinského 9, 812 37 Bratislava, 2024

Holienka, K., Illeová, V., Polakovič, M.: Enhancing  $\gamma$ -Caprolactone Production via a Multienzyme Cascade in *E. coli*: Impact of Oxygen Transfer and Ethanol as a Competitive Inhibitor, Editors: Mihaľ, M., In *51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025*, Jasná, Demänovská Dolina, Slovakia, 2025.

## Enhancing $\epsilon$ -Caprolactone Production via a Multienzyme Cascade in *E. coli*: Impact of Oxygen Transfer and Ethanol as a Competitive Inhibitor

Karol Holienka, Viera Illeová, Milan Polakovič

*Department of Chemical and Environmental Engineering, Faculty of Chemical and Food  
Technology, Slovak University of Technology, Bratislava, Slovakia*

*e-mail: xholienka@gmail.com*

**Key words:**  $\epsilon$ -Caprolactone, Baeyer-Villiger oxidation, enzyme cascade

$\epsilon$ -Caprolactone is an important monomer in the industry, which can be polymerized into polycaprolactone. Traditional chemical synthesis of lactones uses Baeyer-Villiger oxidation (BVO) of ketones using peracids, which has disadvantages such as low selectivity and formation of harmful by-products. In this study, we present a sustainable alternative via whole-cell biocatalysis using a three-enzyme cascade (ADH, ER, CHMO) that efficiently converts 2-cyclohexanone-1-ol to  $\epsilon$ -caprolactone.

We investigated the effect of oxygen transfer coefficient (kLa) and substrate concentration (cs) on the efficiency of oxidation of cyclohexanone to  $\epsilon$ -caprolactone. The results indicate a strong dependence of the cascade rate on the correct ratio between kLa and cs, due to the simultaneous metabolic consumption of oxygen by the cells themselves.

We further studied the effect of ethanol as an inhibitor of ADH, which catalyzes not only the first step of the cascade but also the reaction of cyclohexanone to the undesired by-product cyclohexanol. Our study showed that there is an optimal concentration of ethanol that inhibits the side reaction without significantly suppressing the desired ADH activity. These results contribute to a better understanding of kinetic interactions within multi-enzymatic reaction networks and provide a basis for the design of more efficient biocatalytic processes.

**Acknowledgment:** This work was supported by grants from the Slovak Research and Development Agency (Grant number: APVV-20-0272) and the Scientific Grant Agency of the Slovak Republic (Grant number: VEGA 1/0515).