



Slovak Society of Chemical Engineering
Institute of Chemical and Environmental Engineering
Slovak University of Technology in Bratislava

PROCEEDINGS

51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025

Hotel DRUŽBA
Jasná, Demänovská Dolina, Slovakia
May 27 - 30, 2025

Editors: Assoc. Prof. Mário Mihaľ

ISBN: 978-80-8208-158-2, EAN: 9788082081582

Published by the Faculty of Chemical and Food Technology Slovak Technical University in Bratislava in Slovak Chemistry Library for the Institute of Chemical and Environmental Engineering; Radlinského 9, 812 37 Bratislava, 2024

Koók, L., Rózsenberszki, T., Kovács, S., Kucsera, K.: Production of carboxylic acids from organic waste via electro-fermentation: selectivity, recovery and electrochemical upgrading opportunities, Editors: Mihaľ, M., In *51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025*, Jasná, Demänovská Dolina, Slovakia, 2025.

Production of carboxylic acids from organic waste via electro-fermentation: selectivity, recovery and electrochemical upgrading opportunities

László Koók¹, Tamás Rózsenberszki¹, Szilveszter Kovács¹, Katalin Kucsera¹

¹*Research Group on Bioengineering, Membrane Technology and Energetics, University of Pannonia, Egyetem str. 10. Veszprém, 8200, Hungary*

e-mail: kook.laszlo@mk.uni-pannon.hu

Key words: carboxylic acid, dark fermentation, electro-fermentation, oxidative decarboxylation

Transforming organic waste into useful recoverable products is a challenging issue, which requires highly efficient and integrated technological solutions to obtain value-added resources. In this work, the organic waste-based production of carboxylic acids (CAs) via electrochemically assisted dark fermentation was studied. Compared to conventional dark fermentation, electro-fermentation applies an external reducing power to achieve ‘unbalanced fermentation’, resulting mainly in shifts in product selectivity, yields and kinetics. During the treatment of kitchen waste, electro-fermentation proved to be advantageous in terms of improving chain elongation to caproic acid (C6) at higher organic loads. In addition, the lag-phase during electro-fermentation was significantly reduced compared to the conventional dark fermentation. Microbial consortia analysis of the bulk revealed that although higher caproic acid production occurred during electro-fermentation, C6-related bacteria were present with lower relative abundance, indicating remarkable fermentation efficiency improvement, assumably together with lower biomass yield. The produced CAs – after separation and concentration by e.g. (bipolar) electro-dialysis – can be directly used, or further upgraded to higher-value components. Recently, electrochemical methods gained attention, since those fit well to the trends of electrifying biotechnology. This study assessed the potential of electrochemical upgrading via anodic decarboxylation and dimerization. In this process, the decarboxylated CA radicals can form dimers, resulting in alkanes – e.g. decane from caproic acid (C6) -, or could undergo other pathways, forming esters, alcohols, short alkenes, etc. Using model solutions of pure CAs and mixtures, mainly alkanes and esters were obtained, with increased selectivity to alkanes with longer chain CAs.

In summary, it was shown that both the fermentation, separation and upgrading of the CAs can be improved by using electrochemical approaches, thus allowing efficient waste-to-product processes. This work was supported by the 2024-2.1.1-EKÖP University Research Fellowship Program of the Ministry of Culture and Innovation from the source of the National Research Development and Innovation Fund.