



Slovak Society of Chemical Engineering
Institute of Chemical and Environmental Engineering
Slovak University of Technology in Bratislava

PROCEEDINGS

51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025

Hotel DRUŽBA
Jasná, Demänovská Dolina, Slovakia
May 27 - 30, 2025

Editors: Assoc. Prof. Mário Mihaľ

ISBN: 978-80-8208-158-2, EAN: 9788082081582

Published by the Faculty of Chemical and Food Technology Slovak Technical University in Bratislava in Slovak Chemistry Library for the Institute of Chemical and Environmental Engineering; Radlinského 9, 812 37 Bratislava, 2024

Mihaľ, M., Červeňanský, I., Markoš, J.: Removal of organic acids from fermentation broth by pertraction hollow fiber membrane module, Editors: Mihaľ, M., In *51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025*, Jasná, Demänovská Dolina, Slovakia, 2025.

Removal of organic acids from fermentation broth by pertraction hollow fiber membrane module

Mário Mihal', Ivan Červeňanský, Jozef Markoš

*Institute of Chemical and Environmental Engineering, Faculty of Chemical and Food Technology,
Slovak University of Technology, Radlinského 9, 812 37 Bratislava, Slovakia*

e-mail: mario.mihal@stuba.sk

Key words: pertraction, hollow fiber membrane module, succinic acid, trioctylamine, mathematical modelling

Organic acids are a class of chemical compounds characterized by a carboxyl group and possess acidic properties. They are typically weak acids and are often involved in various biological and industrial processes. Succinic acid, also known as butanedioic acid, is a naturally occurring dicarboxylic organic acid and it is produced by some organisms during anaerobic fermentation and can be found in both plant and animal tissues. Nowadays, as a key platform chemical it is widely used in the chemical, cosmetic, pharmaceutical or food industries. An environmentally friendly alternative to its petrochemical production is the application of microbial fermentation using bacteria such as *Actinobacillus succinogenes* or *Escherichia coli*, which has the potential to reduce its production costs and decrease the environmental impact as they can utilize renewable sources like lignocellulosic biomass, waste streams, or even glycerol. Separation from the fermentation medium and its purification to high purity is also an important part of its biological production as it is often produced together with other organic acids as formic acid, acetic acid or lactic acid. In the process of separation it is beneficial to use several membrane separation techniques such as microfiltration, pertraction, membrane crystallization.

In this work a set of equilibrium experiments in system organic acid/water/organic solvent was performed to give us the information about partition coefficients of separated organic acids. Several pertraction experiments were realized in pertraction unit equipped with hollow fiber membrane contactor Liqui-Cel with the cross flow between shell and fibers. In this system pertraction kinetics of succinic acid or its mixture with other organic acids from the model fermentation medium to the water stripping phase were measured. As immobilized liquid membrane in the pores of the fibers a 20 % mass trioctylamine in octanol was used. To better understand the process of pertraction of organic acids from aqueous phase the individual pertraction experiments were performed at different pH values of the feed and stripping phase and compared also with the prediction of mathematical model.

Acknowledgement: This work was supported by the Slovak Scientific Agency, Grant No. VEGA 1/0658/24, and the Slovak Research and Development Agency under the contract No. APVV-22-0038 and VV-MVP-24-0365.