



Slovak Society of Chemical Engineering  
Institute of Chemical and Environmental Engineering  
Slovak University of Technology in Bratislava

## PROCEEDINGS

51<sup>st</sup> International Conference of the Slovak Society of Chemical Engineering SSCHE 2025

Hotel DRUŽBA  
Jasná, Demänovská Dolina, Slovakia  
May 27 - 30, 2025

Editors: Assoc. Prof. Mário Mihaľ

ISBN: 978-80-8208-158-2, EAN: 9788082081582

Published by the Faculty of Chemical and Food Technology Slovak Technical University in Bratislava in Slovak Chemistry Library for the Institute of Chemical and Environmental Engineering; Radlinského 9, 812 37 Bratislava, 2024

Kotuličová, S., Molnár, T., Majerová, P., Kováč, A., Polakovič, M.: Development and optimization of chromatographic methods for the separation of recombinant tau proteins, Editors: Mihaľ, M., In *51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025*, Jasná, Demänovská Dolina, Slovakia, 2025.

## Development and optimization of chromatographic methods for the separation of recombinant tau proteins

S. Kotuličová<sup>1</sup>, T. Molnár<sup>1</sup>, P. Majerová<sup>2</sup>, A. Kováč<sup>2</sup>, M. Polakovič<sup>1</sup>

<sup>1</sup>*Slovak University of Technology in Bratislava, Faculty of Chemical and Food Technology, Radlinského 9, 812 37 Bratislava*

<sup>2</sup>*Slovak Academy of Sciences, Institute of Neuroimmunology, Dúbravská cesta 9, 845 10 Bratislava 45*

[simona.kotulicova@stuba.sk](mailto:simona.kotulicova@stuba.sk)

**Key words:** Tau proteins, separation, size-exclusion chromatography, hydrophobic interaction chromatography

In neurons, tau proteins play key role in the stability of microtubules and in the efficiency of axonal transport through regulated interactions with tubulins, which are affected by isoform variations and phosphorylation. However, pathological modifications, specifically hyperphosphorylation, lead to microtubule disruption and to the formation of tau aggregates, including oligomers and neurofibrillary tangles. These aggregates negatively impact neuronal function and contribute to neurodegenerative diseases. The intercellular transfer of these aberrant tau proteins induces neuronal cell death and the advancement of neurodegenerative diseases, particularly in Alzheimer's disease.

The objective of this study was to design an effective chromatographic process for the separation of tau protein isoforms. The inherent similarity in primary structure among these isoforms complicates separation due to their comparable physicochemical characteristics. The potential of size-exclusion chromatography, which enables separation within a narrow size range, was explored. Additionally, hydrophobic interaction chromatography was investigated to leverage differences in hydrophobicity among isoforms.

To enhance separation efficiency, size-exclusion chromatography columns with varying pore sizes, connected in series with increasing pore diameter, were tested. Buffers of higher ionic strength were employed to eliminate non-specific interactions. Some isoforms, despite their minor molecular weight differences, were successfully separated using a 0.01M Tris buffer (pH 7) supplemented with 0.5M NaCl and GuHCl as mobile phase.

Initial experiments using hydrophobic interaction chromatography were conducted with a reversed-phase column and a 5-80% acetonitrile gradient in 0.1% formic acid as mobile phase. Preliminary chromatograms showed visible peaks, suggesting potential separation of tau protein isoforms.

### **Acknowledgements**

*This work was supported by grant from the Slovak Research and Development Agency (Grant number: APVV-21-0321).*