



Slovak Society of Chemical Engineering
Institute of Chemical and Environmental Engineering
Slovak University of Technology in Bratislava

PROCEEDINGS

51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025

Hotel DRUŽBA
Jasná, Demänovská Dolina, Slovakia
May 27 - 30, 2025

Editors: Assoc. Prof. Mário Mihaľ

ISBN: 978-80-8208-158-2, EAN: 9788082081582

Published by the Faculty of Chemical and Food Technology Slovak Technical University in Bratislava in Slovak Chemistry Library for the Institute of Chemical and Environmental Engineering; Radlinského 9, 812 37 Bratislava, 2024

Balouch, M., Storchmannová, K., Berka, K., Štěpánek, F.: Drug permeability: commonly misinterpreted bioavailability governing factor, Editors: Mihaľ, M., In *51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025*, Jasná, Demänovská Dolina, Slovakia, 2025.

Drug permeability: commonly misinterpreted bioavailability governing factor

Martin Balouch^{1,2}, Kateřina Storchmannová³, Karel Berka³, František Štěpánek¹

¹*Department of Chemical Engineering, University of Chemistry and Technology Prague, Czech Republic*

²*Zentiva, k.s., U Kabelovny 130, Prague 10 102 37, Czech Republic*

³*Department of Physical Chemistry, Faculty of Science, Palacký University Olomouc, Czech Republic*

e-mail: balouchm@vscht.cz

Key words: permeability, bioavailability, meta-analysis

Permeability is a crucial molecular property in drug discovery that influences pharmacokinetics when drugs cross phospholipid bilayers, such as cell membranes, the gastrointestinal tract, or the blood-brain barrier. Numerous methods exist for determining permeability, including cell line assays (CACO-2, MDCK), cell-free systems like parallel artificial membrane permeability assay (PAMPA), black lipid membrane (BLM), or liposome-based assays. Various *in silico* approaches have also been developed.

We have analyzed already published permeability data (MolMeDB and ChEMBL databases) evaluated four experimental and two computational methods. Results showed poor repeatability even within the same method. For PAMPA, intrinsic and apparent permeabilities can differ by orders of magnitude, suggesting caution when using literature data.

The unstirred water layer permeability (permeability of drug through stationary water layer on the surface of the membrane) causes discrepancies between methods and is often cause of misinterpretation.

Furthermore, we have identified the limitations of studied methods and transferred the hardly imaginable permeability values into real life drug permeation examples which can serve community in order to better interpret both own and literature data. In conclusion, caution is recommended when interpreting or combining permeability data.