

Slovak Society of Chemical Engineering Institute of Chemical and Environmental Engineering Slovak University of Technology in Bratislava

PROCEEDINGS

 $51^{\rm st}$ International Conference of the Slovak Society of Chemical Engineering SSCHE 2025

Hotel DRUŽBA Jasná, Demänovská Dolina, Slovakia May 27 - 30, 2025

Editors: Assoc. Prof. Mário Mihaľ

ISBN: 978-80-8208-158-2, EAN: 9788082081582

Published by the Faculty of Chemical and Food Technology Slovak Technical University in Bratislava in Slovak Chemistry Library for the Institute of Chemical and Environmental Engineering; Radlinského 9, 812 37 Bratislava, 2024

Brzozowski, B., Adamczak, M.: The application of -D- galactosidase from Bifidobacterium bifidum in the synthesis of galactooligosaccharides, Editors: Mihaľ, M., In 51st International Conference of the Slovak Society of Chemical Engineering SSCHE 2025, Jasná, Demänovská Dolina, Slovakia, 2025.

The application of β-D- galactosidase from *Bifidobacterium bifidum* in the synthesis of galactooligosaccharides

Bartosz Brzozowski, Marek Adamczak*

Department of Process Engineering, Equipment and Food Biotechnology, University of Warmia and Mazury in Olsztyn, Jan Heweliusz St. 1, 10-718 Olsztyn, Poland

e-mail: marek.adamczak@uwm.edu.pl

Key words: deep eutectic solvents, galactooligosaccharides, galactosidase, hydrolysis, transgalactosylation

Galactooligosaccharides (GOS) are a class of functional oligosaccharides considered prebiotic. GOS selectively stimulate the growth of beneficial bifidobacteria in the gut, contributing to improved gut health and overall well-being. GOS have been shown to increase probiotic viability and antioxidant activity in fermented products.

The aim of the experiments was to evaluate the preference of β -D-galactosidase from *Bifidobacterium bifidum* in the galactooligosaccharide synthesis reaction. Reactions were carried out in aqueous solutions of lactose and in the environment of deep eutectic liquids (DES). The composition of the reaction mixture was analyzed using liquid chromatography with refractometer and scattered light detector. Favorable synthesis of GOS (about 30%) was found in the reaction catalyzed by NURICA in 20-40% of lactose solutions and at 40-60°C.

At temperatures up to 30°C, the transgalactosylation reaction occurred slowly and less efficiently. During transgalactosylation reactions carried out at 70 and 80°C, the enzyme is inactivated.

In the reaction catalyzed by NURICA in a deeply eutectic liquid medium (choline chloride:glucose, 1:2 M), preferential catalysis of the hydrolysis reaction was observed.

Project funded under the designated subsidy of the Minister of Science and Higher Education Republic of Poland, task entitled: The Research Network of Life Sciences Universities for the Development of the Polish Dairy Industry – Research Project.